

Is small scale convection responsible for the formation of thick igneous crust along volcanic passive margins?

Abdolreza Ghods

Institute for Advanced Studies in Basic Sciences, Zanjan, Iran

Received 20 November 2001; revised 18 March 2002; accepted 3 April 2002; published 16 May 2002.

[1] *Boutillier and Keen* [1999] showed that small scale convection may produce the observed thick igneous crust along volcanic passive margins provided that a non-linear rheology is adopted and the viscosity of the upper mantle is increased suddenly after 132 km of continental crust extension. The sudden viscosity increase is justified by the melt induced dewatering of the upper mantle rocks. Through this comparative study, it is shown that inclusion of the pressure of the newly formed igneous crust into the melting model, and/or adopting a more realistic viscosity law, significantly reduces the thickness of igneous crust produced by Boutillier and Keen's models. Furthermore, it is shown that melt migration would narrow the width of the thick igneous crust. Results of this study show that small scale convection may not play a significant role in the formation of the observed thick igneous crust along volcanic passive margins.

INDEX TERMS: 8109 Tectonophysics: Continental tectonics—extensional (0905); 8159 Evolution of the Earth: Rheology—crust and lithosphere; 8120 Tectonophysics: Dynamics of lithosphere and mantle—general; 3230 Mathematical Geophysics: Numerical solutions; 3220 Mathematical Geophysics: Nonlinear dynamics

1. Introduction

[2] Evolution of continental rifting to sea-floor spreading in volcanic passive margins is accompanied by formation of thick igneous rocks within a narrow ocean-continent transition zone (50–100 km). Unusually thick igneous crust of 20–30 km has been detected beneath the narrow ocean-continent transition zone of volcanic passive margins [e.g., *White and McKenzie*, 1989; *White*, 1992]. Northern North Atlantic and U.S. East Coast Margin are among the typical examples of volcanic passive margins where during a short period of 2–3 m.y. over a few million km³ of igneous rocks was produced [*White et al.*, 1987; *Holbrook*, 1993]. The sheer volume of igneous rocks rivals that of continental flood basalts and thus is considered to be in the category of the large igneous provinces [*Coffin and Eldholm*, 1994].

[3] During the last two decades, there has been many efforts to explain the observation of the thick igneous crust along volcanic passive margins. One school of thought considers an anomalously hot upper mantle as a responsible mechanism for formation of the thick igneous crust. The source of such temperature anomaly is postulated to be either related to the presence of hot mantle plumes [*White and McKenzie*, 1989; *White*, 1988] or the thermal gradient created by insulating effect of the large continents [*Anderson*, 1982, 1992; *Gurnis*, 1988]. The other school of thought holds the small scale convection happening at the initial stages of continental rifting responsible for the formation of the thick igneous crust [*Mutter and Wernicke*, 1988; *Zehnder et al.*, 1990]. *Mutter and Wernicke* [1988] hypothesized that the initial sharp thermal contrast between the rifting thinned continent and the asthenosphere induces small scale convection that increases the flux of material

crossing the solidus and in turn increases the total volume of melt. By progress of the sea-floor spreading and cooling of the rifted continent, the sharp thermal contrast and its associated small scale convection fades away after initiation of sea-floor spreading, and thus a normal oceanic crust forms afterward. Recently, *Boutillier and Keen* [1999] (hereafter called B&K) tested the small-scale convection scenario and showed that considering a non-linear rheology leads to a small scale convection which successfully produces the observed thick igneous crust with a width close to that of observed. In their successful model (e.g., following the authors I call the successful model the wet model) the non-linear rheology also depends on melting. The viscosity of the 80 km of the upper mantle is increased suddenly after 132 km of continental crust extension as an attempt to simulate the melt induced viscosity increase of the upper mantle rocks. The authors ignore the effect of melt depletion buoyancy and pressure of the newly formed igneous crust. In computing the thickness of igneous crust, the authors assume that melt is instantly underplated directly above where it is formed.

[4] In this paper, I will report the results of a comparative study of my models with B&K's. By investigating the effect of the melt depletion buoyancy, the pressure of newly formed igneous crust and a more realistic melt-dependent rheological model on the volume of melt production, I will show that some of the assumptions used in B&K's models are unrealistic. As it will be shown later, considering melt depletion buoyancy for the upper mantle rocks will not significantly change the results obtained by the authors. However, including the effect of the pressure of the newly formed igneous crust and/or adopting of a more realistic viscosity law will significantly reduces the thickness of the igneous crust. The new viscosity law simulates the melt induced viscosity increase of the solid upper mantle rocks by a gradual increase of viscosity upon progress of melting. Finally, I show that ignoring melt migration in the partially molten upper mantle plays an important role in redistributing melt production to a wider area and thus avoiding creation of a localized narrow thick oceanic crust in the B&K wet model.

2. Numerical Model

[5] I model the corner flow induced by continental rifting in a two dimensional (2D) computational frame of 2100 km wide and 700 km deep. My base model has the same settings of the B&K wet model. The top boundary moves horizontally with a specified spreading velocity of 5 km/m.y. in all of my models. The bottom boundary is free-slip. Mass is allowed to only enter/exit horizontally from the right boundary. Temperature is fixed at the top and bottom boundaries to zero and 1350°C, respectively. The initially assigned temperature field changes linearly within the continental lithosphere and is constant at depths beneath. Viscosity depends on strain rate ($\dot{\epsilon}$), temperature (T) and pressure (P) according to following equation of state,

$$\eta_s = A \dot{\epsilon}_r^{(1-n)/n} \exp\left(\frac{E + PV}{R(T + 273)}\right) \quad (1)$$

Table 1 Physical Parameters Used in this Paper

Parameter	Value
Mantle density (ρ_p^o)	3300 kg m ⁻³
Reference shear viscosity (η_o)	1.29×10^{19} Pa s
Viscosity activation energy (E)	444 kJ/mol
Viscosity activation volume (V)	15×10^{-6} m ³ /mol
Viscosity strain rate component (n)	3.356
Thermal diffusivity (κ)	8.0×10^{-7} m ² /s
Latent heat of melting (H)	1.2×10^9 J m ⁻³
Thermal expansion coefficient (α)	3.2×10^{-5} °C ⁻¹
Melting expansion coefficient (β)	0.024 ^a

^a Sparks and Parmentier, 1994.

values of E , V and n are the same as those B&K' models (Table 1). A reference strain rate ($\dot{\epsilon}_r$) of 1.67×10^{-17} and a reference viscosity of 3.6×10^{23} are assumed at 700 km depth, the bottom of my computational domain, to calculate value of A . In order to simulate the effect of melt induced dewatering on viscosity, B&K increased viscosity by a factor of 140 at depths above 80 km when 132 km of lateral extension is completed. Following B&K, the initial thickness of continental lithosphere is reduced by an extension factor of five over a narrow region of 16 km at the rift axis. This is an attempt to simulate a sharp initial necking of continental lithosphere.

[6] An iterative finite volume method [Patankar, 1980] is adopted to solve equations of mass, momentum and energy. Following B&K, the melt phase is completely ignored in the governing equations. The numerical results presented in this paper are calculated in a non-uniform rectangular grid with 106×81 nodes with high resolution for an area of 87.5 km depth from the surface and 262 km width from the ridge axis, where melting usually occurs. Vertical grid-spacing within the high resolution area is 1.8 km and varies between about 6 to 27 km outside of the region. Horizontal grid-spacing within the high-resolution area varies between 1.5 to 4.2 km and increases from 16 km to a maximum of 88 km outside the high-resolution area. The smooth horizontal and vertical gradients of velocity and temperature fields at the edge and outside of the high-resolution area justifies the use of the rather large grid-spacings within the low resolution area.

3. Melting Model

[7] I use a fractional melting model [Ghods and Arkani-Hamed, 2000] where melt is extracted completely. The fractional melting model is a modified version of McKenzie and Bickle's [1988] batch melting model and is constructed by replacing the liquidus temperature (T_l^o) of peridotite by the liquidus temperature of pure forsterite (i.e., $T_l^o = 2164 + 4.77 P_H$, [Presnall and Walter, 1993]). Furthermore, to account for the dynamic nature of melting, the depletion field \mathcal{M} is advected in each time step using the following pure advective transport equation,

$$\frac{\partial \mathcal{M}}{\partial t} + V \cdot \nabla \mathcal{M} = 0 \quad (2)$$

where V is velocity of the upper mantle rocks. \mathcal{M} quantifies the depletion of a partially molten rock and indicates the ratio of the pristine rock that should melt to produce the solid residue. The above equation is solved using monotonic second-order upwind method (Sweby, 1984). The fraction of melt produced in a given time step $\Delta \mathcal{M}$ is computed using the following coupled equations,

$$\begin{aligned} \rho_s C_P T_1 &= \rho_s C_P T_2 + \Delta \mathcal{M} H, \\ \Delta \mathcal{M} &= \frac{T_2 - T_s}{T_l^o - T_s} (1 - \mathcal{M}), \end{aligned} \quad (3)$$

By solving the above equations we can also calculate the temperature field corrected for the absorption of the latent heat of melting, T_2 , and update the depletion field \mathcal{M} . T_1 is temperature before the correction. The value of latent heat of melting H is given in Table 1. T_s is the instantaneous solidus temperature of the solid residue at the end of the previous time step. To account for the increase of T_s through melting, T_s is updated at the end of each time step using $\mathcal{M} = \frac{T - T_s^o}{T_l^o - T_s^o}$ where T_s^o is solidus temperature of pristine dry peridotite, and then setting $T_s = T$. The removal of the melt in a fractional melting model increases T_s because the residue becomes more refractory.

[8] There are two minor differences between B&K's melting model and that of mine. First, I correct temperature field to account for the latent heat of melting whereas B&K do not. Second, my models use a fractional melting model which corrects the original batch melting model of McKenzie and Bickle [1988] to take into account the effect of fractional melting. This has been accomplished by considering a liquidus curve whose temperature is generally larger than the batch melting liquidus curve used by B&K. A comparative study of the two melting models shows that the above mentioned differences have very slight effects on the results obtained by B&K's models. Therefore I use my fractional melting model when reproducing B&K' models.

4. Modeling

[9] In the first set of experiments, the effect of inclusion of melt depletion buoyancy into the B&K wet model is investigated. Addition of melt depletion buoyancy to mid-ocean ridge models increases melt production at very slow spreading rates (e.g., Cordery and Phipps Morgan, 1993), however melt depletion buoyancy would decrease the rate of recycling of the depleted upper mantle rocks into the melting zone of the upper mantle. Inclusion of melt buoyancy increases the peak crustal thickness by about 2 km (Figure 1). However, such increase of crustal thickness once compared with the observed thick igneous crust is marginal. This implies that the melt buoyancy can be safely ignored as was done in B&K's models.

[10] In the second set of experiments, the effect of the thickness of newly formed igneous rocks on the volume of melt production at later stages of rifting is investigated. Volcanism and underplating at early stages of rifting forms a new igneous layer at the edge of a rifting crust. Such a layer exerts extra pressure on the melt producing zone and thus may significantly reduce the thickness




Figure 1. Crustal thickness versus time. The dashed line is the B&K wet model and the solid line is the model that consider melt depletion buoyancy. The horizontal axis is the time from the initiation of rifting.




Figure 2. Crustal thickness versus time for the B&K wet model (the dashed line) and the modified B&K's model which consider the pressure of the newly formed igneous crust (the solid line).

of the melting zone and the total melt volume produced off the rift axis at later stages of rifting. To quantify the effect of pressure of the overlying igneous layer on the melt production, I include the pressure of the overlying igneous crust in the melting model. Figure 2 compares the crustal thickness produced by the later model and that of the B&K wet model. It shows that the crustal thickness decreases significantly by the addition of the pressure of the newly formed igneous crust.

[11] In the third set of experiments, the effect of melt-induced viscosity increase on the crustal thickness is investigated using a sudden and a gradual viscosity increase. B&K increase the viscosity of the top 80 km of the upper mantle by a factor of 140 after 132 km of crustal extension in order to simulate the effect of melt induced dewatering of the upper mantle. However, the viscosity of the upper mantle increases gradually in response to the gradual melt induced dewatering of the upper mantle [Hirth and Kohlstedt, 1996]. In order to show that the sudden increase of viscosity is very critical in the success of B&K's models, I run a model in which viscosity increases linearly with the increase of melt percentage and reaches to a maximum of 140 times than that obtained by equation (1) at melt fraction of 15% and above. Figure 3 compares crustal production of the two models. It shows that the peak crustal




Figure 3. Crustal thickness versus time for the B&K wet model (the dashed line) and the modified B&K's model which increases viscosity linearly by increase of melt percentage (the solid line).




Figure 4. Crustal thickness versus time for the B&K wet model (the dashed line) and the modified B&K's model which instantly delivers melt beneath the ridge axis (the solid line).

thickness for my model (the solid line) is much smaller than that of B&K's model (the dashed line). This result implies that small scale convection may not play a significant role in the formation of the thick igneous crust observed along volcanic passive margins. Adopting a viscosity law which allow for larger increase of viscosity at earlier stages of melting would increase viscosity of the upper mantle sooner and thus causes a more pronounced reduction of the peak crustal thickness.

[12] In the last set of experiments, the combined effect of the thickness of newly formed igneous rocks and melt induced viscosity increase on the crustal thickness is investigated. The model with both effects included produces a smaller peak of crustal thickness (about 5 km) than that which only includes the effect of melt induced viscosity increase (6 km).

5. Discussion and Conclusions

[13] Through this comparative study, it is shown that small scale convection can not reproduce the observed crustal thickness along volcanic passive margins when a more realistic melting model and/or viscosity law are used. Crustal production produced by the B&K wet model is significantly decreased when the effect of the pressure of overlying newly igneous crust is included and a viscosity law which increases viscosity linearly by increase of melt percentage is adopted. It is also shown that B&K results are not sensitive to the melt depletion buoyancy.

[14] Using a two-phase flow model, Ghods and Arkani-Hamed [2000] showed that half of melt generated beneath the mid-ocean ridge delivers itself beneath the ridge axis and the rest solidifies beneath the lithosphere. In their model, they consider a completely reversible melting-freezing model which certainly gives an upper limit to the extent of freezing in their model. Ghods and Arkani-Hamed's [2000] results suggest that melt migration within the partially molten upper mantle beneath mid-ocean ridge effectively focuses melt toward the ridge axis. The critical components in focusing melt beneath mid-ocean ridge are the sloping impermeable base of the lithosphere and the compressible nature of a partially molten zone of the upper mantle. Both components are present beneath a rift axis and furthermore the slope of the lower boundary of the rifting lithosphere is even larger. Therefore, melt migration would focus melt effectively beneath the rift axis. This focusing effect is not considered in the B&K's models. Figure 4 compares crustal production produced by the B&K wet model and the one which simulate melt migration by instant delivery of melt beneath the ridge axis. The maximum crustal thickness of the

model with melt migration is significantly (i.e., by more than 4 km) larger than that without it, however, the width of the thick crust is narrower by a factor of about two than that of the B&K wet model and that of the observed. This result still provides us a further reason to not hold the small scale convection as a responsible mechanism for the creation of the thick igneous crust produced along volcanic passive margins.

[15] **Acknowledgments.** This research was supported by Institute for advanced studies in basic sciences (IASBS), Zanjan, Iran. I would like to thank Farhad Sobouti (Dep. of Physics, IASBS) for his constructive comments. I also like to thank one anonymous reviewer for his/her very useful comments.

References

- Anderson, D. L., Hotspots, polar wander, Mesozoic convection and the geoid, *Nature*, 297, 391–393, 1982.
- Anderson, D. L., Large igneous provinces: Non-plume events, *Trans. AGU*, 73(43), 531, 1992. Fall Meet. suppl.
- Boutillier, R. R., and C. E. Keen, Small-scale convection and divergent plate boundaries, *J. Geophys. Res.*, 104, 7389–7403, 1999.
- Coffin, M. F., and O. Eldholm, Large igneous provinces: Crustal structure, dimension, and external consequences, *Rev. Geophys.*, 32, 1–36, 1994.
- Cordery, M. J., and J. Phipps Morgan, Convection and modeling at mid-ocean ridges, *J. Geophys. Res.*, 98, 19,477–19,503, 1993.
- Ghods, A., and J. Arkani-Hamed, Melt migration beneath mid-ocean ridges, *Geophys. J. Int.*, 140, 687–697, 2000.
- Gurnis, M., Large scale mantle convection and the aggregation and dispersal of supercontinents, *Nature*, 332, 695–699, 1988.
- Hirth, G., and D. L. Kohlstedt, Water in oceanic upper mantle: Implications of rheology, melt extraction and the evolution of the lithosphere, *Earth Planet Sci. Lett.*, 144, 93–108, 1996.
- Holbrook, P. B., Kelemen, Large igneous province on the US Atlantic margin and implications for magmatism during continental breakup, *Nature*, 364, 433–436, 1993.
- McKenzie, D., and M. J. Bickle, The volume and composition of melt generated by extension of the lithosphere, *J. Petrol.*, 29, 625–679, 1988.
- Mutter, J. C., and B. Wernicke, Comments on rifting and passive margins evolution in light of some recent studies, National Academy Press, pp. 208–229, Washington, D.C., 1988.
- Patankar, S. V., *Numerical heat transfer and fluid flow*, 329 pp, Hemisphere publishing corporation, 1980.
- Presnall, D. C., and M. J. Walter, Melting of forsterite, Mg_2SiO_4 , from 9.7 to 16.5 GPa, *J. Geophys. Res.*, 98, 19,777–19,783, 1993.
- Sparks, D. W., and E. M. Parmentier, The generation and migration of partial melt beneath oceanic spreading centers, in *Magmatic System*, edited by M. P. Ryan, pp. 55–76, Academic Press, San Diego, 1994.
- White, R. S., A hot-spot model for early Tertiary volcanism in the N. Atlantic, in *Early Tertiary Volcanism and the Opening of the NE Atlantic*, edited by A. C. Morton and L. M. Parson, *Geol. Soc. London Spec. Pub.*, 39, 3–13, 1988.
- White, R. S., Crustal structure and magmatism of North Atlantic continental margins, *J. Geophys. Res.*, 149, 841–854, 1992.
- White, R. S., and D. P. McKenzie, Magmatism at rift zones: The generation of volcanic continental margins and flood basalts, *J. Geophys. Res.*, 94, 7685–7729, 1989.
- White, R. S., G. D. Spence, D. P. McKenzie, G. K. Westbrook, and A. N. Bowen, Magmatism at rifted continental margins, *Nature*, 330, 439–444, 1987.
- Zehnder, C. M., J. C. Mutter, and P. Buhl, Deep seismic and geochemical constraints on the nature of rift-induced magmatism during breakup of the North Atlantic, *Tectonophysics*, 173, 545–565, 1990.

A. Ghods, Institute for Advanced Studies in Basic Sciences, Zanjan, Iran.